

**NEET-UG – 2013 TEST PAPER WITH SOLUTIONS  
(HELD ON SUNDAY 05<sup>th</sup> MAY, 2013)**

**136.** The value of Planck's constant is  $6.63 \times 10^{-34}$  Js. The speed of light is  $3 \times 10^{17}$  nm s<sup>-1</sup>. Which value is closest to the wavelength in nanometer of a quantum of light with frequency of  $6 \times 10^{15}$  s<sup>-1</sup> ?

- (1) 75 (2) 10  
(3) 25 (4) 50

**Ans. (4)**


**Sol.**  $\lambda = \frac{c}{\nu} = \frac{3 \times 10^{17} \text{ nm s}^{-1}}{6 \times 10^{15} \text{ s}^{-1}} = 50 \text{ nm}$

**137.** The radical,  is aromatic because it

has :-

- (1) 6p-orbitals and 7 unpaired electrons  
(2) 6p-orbitals and 6 unpaired electrons  
(3) 7p-orbitals and 6 unpaired electrons  
(4) 7p-orbitals and 7 unpaired electrons

**Ans. (2)**

**Sol.** , Benzyl free radical is aromatic as


per Huckel's rule it has  $6\pi$  electrons present in p-orbital of carbon atoms involved in formation of benzene ring (Aromatic nature).



**138.** Which of the following is electron-deficient ?

- (1) PH<sub>3</sub> (2) (CH<sub>3</sub>)<sub>2</sub>  
(3) (SiH<sub>3</sub>)<sub>2</sub> (4) (BH<sub>3</sub>)<sub>2</sub>

**Ans. (4)**



**139.** Which of the following statements about the interstitial compounds is **incorrect** ?

- (1) They have higher melting points than the pure metal  
(2) They retain metallic conductivity  
(3) They are chemically reactive  
(4) They are much harder than the pure metal

**Ans. (3)**

**140.** How many grams of concentrated nitric acid solution should be used to prepare 250 mL of 2.0 M HNO<sub>3</sub>?

- (1) 54.0 conc. HNO<sub>3</sub>  
(2) 45.0 conc. HNO<sub>3</sub>  
(3) 90.0 conc. HNO<sub>3</sub>  
(4) 70.0 conc. HNO<sub>3</sub>

**Ans. (2)**

**Sol.**  $M = \frac{W \times 1000}{M_w \times V_{\text{solution}} (\text{mL})}$

$2 = \frac{W \times 1000}{63 \times 250}$

$W = 31.5 \text{ g}$

70% HNO<sub>3</sub> means 70g HNO<sub>3</sub> is present in 100g solution.

$\therefore 31.5 \text{g HNO}_3 \text{ will be present in } \frac{100}{70} \times 31.5$   
 $= 45 \text{ g of solution}$

**141.** Which of the following lanthanoid ions is diamagnetic?

(Atoms, Ce = 58, Sm = 62, Yb = 70)

- (1) Yb<sup>2+</sup> (2) Ce<sup>2+</sup>  
(3) Sm<sup>2+</sup> (4) Eu<sup>2+</sup>

**Ans. (1)**

**Sol.** diamagnetic – no w.e<sup>-</sup> present

Yb<sup>2+</sup> = 70 [Xe] 6s<sup>0</sup> 4f<sup>14</sup> 5d<sup>0</sup> U.e<sup>-</sup> = zero.

**142.** Which one of the following molecules contains no  $\pi$  bond ?

- (1) NO<sub>2</sub> (2) CO<sub>2</sub>  
(3) H<sub>2</sub>O (4) SO<sub>2</sub>

**Ans. (3)**


143. Based on equation  $E = -2.178 \times 10^{-18} \text{ J} \left( \frac{Z^2}{n^2} \right)$

certain conclusions are written. Which of them is **not** correct ?

- (1) For  $n = 1$ , the electron has a more negative energy than it does for  $n = 6$  which means that the electron is more loosely bound in the smallest allowed orbit.
- (2) The negative sign in equation simply means that the energy of electron bound to the nucleus is lower than it would be if the electrons were at the infinite distance from the nucleus
- (3) Larger the value of  $n$ , the larger is the orbit radius
- (4) Equation can be used to calculate the change in energy when the electron change orbit


Ans. (1)

Sol. Correct answer is - for  $n = 1$  the electron has more negative energy than it does for  $n = 6$  which means that the electron is less loosely bound in the smallest allowed orbit.






- (1)  $\text{H}^+/\text{H}_2\text{O}$
- (2)  $\text{HgSO}_4/\text{H}_2\text{SO}_4$
- (3)  $\text{Cu}_2\text{Cl}_2$
- (4)  $\text{H}_3\text{PO}_2$  and  $\text{H}_2\text{O}$

Ans. (4)




145. The order of stability of the following tautomeric compounds is :-



- (1)  $\text{II} > \text{III} > \text{I}$
- (2)  $\text{I} > \text{II} > \text{III}$
- (3)  $\text{III} > \text{II} > \text{I}$
- (4)  $\text{II} > \text{I} > \text{III}$


Ans. (3)

Sol. Stability order is



Intra molecular H-Bonding and conjugated diene and it has involved more acidic -H[-CH<sub>2</sub>-]

(No conjugated diene)



Involvement of less acidic -H in tautomerism

146. Nylon is an example of :-

- (1) Polythene
- (2) Polyester
- (3) Polysaccharide
- (4) Polyamide

Ans. (4)

Sol. Nylon has polyamide linkage. It is formed by condensation reaction of amines and carboxylic acid groups.

147. XeF<sub>2</sub> is isostructural with :-

- (1) BaCl<sub>2</sub>
- (2) TeF<sub>2</sub>
- (3) ICl<sub>2</sub><sup>-</sup>
- (4) SbCl<sub>3</sub>


Ans. (3)

148. The basic structural unit of silicates is :-

- (1) SiO<sub>4</sub><sup>2-</sup>
- (2) SiO<sup>-</sup>
- (3) SiO<sub>4</sub><sup>4-</sup>
- (4) SiO<sub>3</sub><sup>2-</sup>

Ans. (3)

Sol. Silicates - Salt of silicic acid H<sub>4</sub>SiO<sub>4</sub>






silicate ion

149. Which of the following structure is similar to graphite?


- (1) B<sub>2</sub>H<sub>6</sub>
- (2) BN
- (3) B
- (4) B<sub>4</sub>C

Ans. (2)

150. The structure of isobutyl group in an organic compound is :-



Ans. (2)



**151.** The number of carbon atoms per unit cell of diamond unit cell is :-

- (1) 1      (2) 4      (3) 8      (4) 6

**Ans. (3)**

**Sol.** In the diamond cubic unit cell, there are eight corner atoms, six face centered atoms and four more atoms inside the structure.

$\therefore$  Number of atoms present in a diamond cubic unit cell  
 $= 1 + 3 + 4 = 8$  atoms

**152.** An excess of  $\text{AgNO}_3$  is added to 100 mL of a 0.01 M solution of dichlorotetraaquachromium(III) chloride. The number of moles of  $\text{AgCl}$  precipitated would be :-

- (1) 0.01      (2) 0.001      (3) 0.002      (4) 0.003

**Ans. (2)**

**153.** What is the maximum numbers of electrons that can be associated with the following set of quantum numbers ?

$n = 3, l = 1$  and  $m = -1$

- (1) 2      (2) 10      (3) 6      (4) 4

**Ans. (1)**


**Sol.**  $n = 3, l = 1$  and  $m = -1$  represent a 3p-orbital so maximum 2 electrons can be accommodate in it.

**154.** Which of these is not a monomer for a high molecular mass silicone polymer ?

- (1)  $\text{PbSiCl}_3$       (2)  $\text{MeSiCl}_3$   
 (3)  $\text{Me}_2\text{SiCl}_2$       (4)  $\text{Me}_3\text{SiCl}$

**Ans. (4)**

**Sol.** Silicones are organo-silicon polymer



Since  $\text{Me}_3\text{SiCl}$  contain only one Cl therefore it can't form high molecular mass silicon polymer

It act a chain terminating organo silane

**155.** A reaction having equal energies of activation for forward and reverse reactions has :-

- (1)  $\Delta H = \Delta G = \Delta S = 0$       (2)  $\Delta S = 0$   
 (4)  $\Delta G = 0$       (4)  $\Delta H = 0$

**Ans. (4)**

**Sol.**  $\Delta H = (E_a)_f - (E_a)_b$

Given :  $(E_a)_f = (E_a)_b$

$\therefore \Delta H = 0$


**156.** At  $25^\circ\text{C}$  molar conductance of 0.1 molar aqueous solution of ammonium hydroxide is  $9.54 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$  and at infinite dilution its molar conductance is  $238 \text{ ohm}^{-1} \text{ cm}^2 \text{ mol}^{-1}$ . The degree of ionisation of ammonium hydroxide at the same concentration and temperature is :-

- (1) 40.800%      (2) 2.080%  
 (3) 20.800%      (4) 4.008%


**Ans. (4)**

**Sol.**  $\% \alpha = \frac{\lambda^c}{\lambda^\infty} \times 100 = \frac{9.54}{238} \times 100 = 4.008\%$

**157.** Structure of the compound whose IUPAC name is 3-Ethyl-2-hydroxy-4-methylhex-3-en-5-ynoic acid is :-






**Ans. (3)**




3-Ethyl-2-hydroxy-4-methylhex-3-en-5-ynoic acid

**158.** Among the following ethers, which one will produce methyl alcohol on treatment with hot concentrated HI ?



**Ans. (4)**

**Sol.** C-O bond cleavage of ether depends on mechanism involved in reaction which can be  $\text{SN}^1$  or  $\text{SN}^2$ . If any one group can form stable carbocation like tertiary butyl group then reaction follow's  $\text{SN}^1$  mechanism



**159.** Antiseptics and disinfectants either kill or prevent growth of microorganisms. Identify which of the following statements is **not true** :-

- (1) Disinfectants harm the living tissues
- (2) A 0.2% solution of phenol is an antiseptic while 1% solution acts as a disinfectant
- (3) Chlorine and Iodine are used as strong disinfectants
- (4) Dilute solutions of Boric acid and Hydrogen Peroxide are strong antiseptics

**Ans. (4)**

**160.** A magnetic moment of 1.73 BM will be shown by one among the following :-

- (1)  $[\text{CoCl}_6]^{4-}$
- (2)  $[\text{Cu}(\text{NH}_3)_4]^{2+}$
- (3)  $[\text{Ni}(\text{CN})_4]^{2-}$
- (4)  $\text{TiCl}_4$

**Ans. (2)**

**Sol.** Magnetic moment 1.73 BM


$$\mu = \sqrt{n(n+2)} \text{ B.M}$$

n = no. of unpaired  $e^-$


$$\mu = 1.73$$

$$1.73 = \sqrt{n(n+2)} \text{ B.M}$$

$$n = 1$$




$\text{Cl}^-$  (weak field ligand)  $t_2g^5 e_g^2$  unpaired  $e^- = 3$




$\text{NH}_3$  Strong field ligand, hybridisation  $dsp^2$

\* one  $e^-$  of 3d jumps into 4p subshell.


$$\text{unpaired } e^- = 1$$



$\text{CN}^-$  - Strong field ligand  $dsp^2$



**161.**  $\text{KMnO}_4$  can be prepared from  $\text{K}_2\text{MnO}_4$  as per the reaction :-







The reaction can go to completion by removing  $\text{OH}^-$  ions by additions :-

- (1)  $\text{SO}_2$
- (2)  $\text{HCl}$
- (3)  $\text{KOH}$
- (4)  $\text{CO}_2$


**Ans. (4)**

**162.** Reaction by which Benzaldehyde cannot be prepared :-



**Ans. (1)**





$$E_{\text{O.P.}} = E_{\text{O.P.}}^\circ - \frac{0.059}{n} \log \frac{[\text{H}^+]}{(\text{P}_{\text{H}_2})^{1/2}}$$

$$E_{\text{O.P.}} = 0 - \frac{0.059}{1} \log \frac{10^{-10}}{(1)^{1/2}}$$

$$(\text{pH} = 10, [\text{H}^+] = 10^{-10} \text{ M})$$


$$E_{\text{O.P.}} = 0.59 \text{ V}$$

**170.** Which of the following is a polar molecule ?


- (1)  $\text{XeF}_4$  (2)  $\text{BF}_3$  (3)  $\text{SF}_4$  (4)  $\text{SiF}_4$

**Ans. (3)**



**Sol.** Unsymmetrical distribution of  $\text{e}^-$  cloud leads to the formation of polar molecule



**171.** A button cell used in watches function as following



If half cell potentials are




$$E^\circ = 0.34\text{V}$$



The cell potential will be :-

- (1) 1.34 V (2) 1.10 V (3) 0.42 V (4) 0.84 V

**Ans. (2)**



$$E^\circ = 0.34 \text{ V}$$



$$E_{\text{cell}}^\circ = (E_{\text{R.P.}}^\circ)_{\text{cathode}} - (E_{\text{R.P.}}^\circ)_{\text{anode}}$$

$$E_{\text{cell}}^\circ = 0.34 - (-0.76) = 1.10 \text{ V}$$

$$E_{\text{cell}} = E_{\text{cell}}^\circ = 1.10\text{V}$$

**172.** Which of these is **least likely** to act as a Lewis base?

- (1)  $\text{PF}_3$  (2)  $\text{CO}$  (3)  $\text{F}^-$  (4)  $\text{BF}_3$

**Ans. (4)**

**173.** Which of the following compounds will not undergo Friedal-Craft's reaction easily :-

- (1) Toluene (2) Cumene  
(3) Xylene (4) Nitrobenzene

**Ans. (4)**



**Sol.** Friedal Craft reaction fails when strong deactivating group is attached with benzene ring.

**174.** Which is the monomer of Neoprene in the following ?

- (1)  $\text{CH}_2=\text{CH}-\text{C}=\text{CH}$  (2)  $\text{CH}_2=\text{CH}-\text{CH}=\text{CH}_2$   
(3)  $\text{CH}_2=\underset{\text{CH}_3}{\text{C}}-\text{CH}=\text{CH}_2$  (4)  $\text{CH}_2=\underset{\text{Cl}}{\text{C}}-\text{CH}=\text{CH}_2$

**Ans. (4)**

**Sol.**



**175.**  $6.02 \times 10^{20}$  molecules of urea are present in 100mL of its solution. The concentration of solution is :-

- (1) 0.1 M (2) 0.02 M  
(3) 0.01 M (4) 0.001M

**Ans. (3)**

**Sol.** Number of moles =  $\frac{\text{number of molecules}}{N_A}$

$$= \frac{6.02 \times 10^{20}}{6.02 \times 10^{23}} = 10^{-3} \text{ mol}$$

$$\text{Molar conc.} = \frac{n \times 1000}{V_{\text{solution}} (\text{mL})} = \frac{10^{-3} \times 1000}{100}$$

$$\text{Molar conc.} = 0.01 \text{ M}$$

**176.** Maximum deviation from ideal gas is expected from :

- (1)  $\text{NH}_3(\text{g})$  (2)  $\text{H}_2(\text{g})$  (3)  $\text{N}_2(\text{g})$  (4)  $\text{CH}_4(\text{g})$

**Ans. (1)**


**Sol.**  $\text{NH}_3$  will show maximum deviation from ideal gas due to dipole-dipole attraction.

**177.** Which of the following is paramagnetic ?

- (1)  $\text{NO}^+$  (2)  $\text{CO}$  (3)  $\text{O}_2^-$  (4)  $\text{CN}^-$

**Ans. (3)**

**Sol.**  $\text{O}_2^- \rightarrow 15 e^-$  contains one unpaired  $e^-$  in ABMO.



**178.** Identify the correct order of solubility in aqueous medium :

- (1)  $\text{Na}_2\text{S} > \text{ZnS} > \text{CuS}$   
 (2)  $\text{CuS} > \text{ZnS} > \text{Na}_2\text{S}$   
 (3)  $\text{ZnS} > \text{Na}_2\text{S} > \text{CuS}$   
 (4)  $\text{Na}_2\text{S} < \text{CuS} > \text{ZnS}$

**Ans. (1)**

**179.** What is the activation energy for a reaction if its rate doubles when the temperature is raised from  $20^\circ\text{C}$  to  $35^\circ\text{C}$  ? ( $R = 8.314 \text{ J mol}^{-1} \text{ K}^{-1}$ )

- (1)  $15.1 \text{ kJ mol}^{-1}$  (2)  $342 \text{ kJ mol}^{-1}$   
 (3)  $269 \text{ kJ mol}^{-1}$  (4)  $34.7 \text{ kJ mol}^{-1}$

**Ans. (4)**

**Sol.** 
$$\log \frac{K_2}{K_1} = \frac{E_a}{2.303R} \left( \frac{1}{T_1} - \frac{1}{T_2} \right)$$

$$\frac{r_2}{r_1} = \frac{K_2}{K_1} = 2$$

$$\Rightarrow \log 2 = \frac{E_a}{2.303 \times 8.314 \times 10^{-3}} \left( \frac{1}{293} - \frac{1}{308} \right)$$

$$\Rightarrow E_a = 34.7 \text{ KJ mol}^{-1}$$

**180.** Which is the strongest acid in the following ?

- (1)  $\text{H}_2\text{SO}_3$  (2)  $\text{H}_2\text{SO}_4$  (3)  $\text{HClO}_3$  (4)  $\text{HClO}_4$

**Ans. (4)**