

**MAXIMA & MINIMA**

1. Let  $f(x)$  be a polynomial of degree 5 such that  $x = \pm 1$  are its critical points. If  $\lim_{x \rightarrow 0} \left( 2 + \frac{f(x)}{x^3} \right) = 4$ , then which one of the following is not true?

- (1)  $f$  is an odd function
- (2)  $x = 1$  is a point of minima and  $x = -1$  is a point of maxima of  $f$ .
- (3)  $x = 1$  is a point of maxima and  $x = -1$  is a point of minimum of  $f$ .
- (4)  $f(1) - 4f(-1) = 4$

2. Let  $f(x)$  be a polynomial of degree 3 such that  $f(-1) = 10, f(1) = -6, f(x)$  has a critical point at  $x = -1$  and  $f'(x)$  has a critical point at  $x = 1$ . Then  $f(x)$  has a local minima at  $x =$  \_\_\_\_\_.

3. Let a function  $f : [0, 5] \rightarrow \mathbf{R}$  be continuous,  $f(1) = 3$  and  $F$  be defined as :

$$F(x) = \int_1^x t^2 g(t) dt, \text{ where } g(t) = \int_1^t f(u) du.$$

Then for the function  $F$ , the point  $x = 1$  is :

- (1) a point of local minima.
- (2) not a critical point.
- (3) a point of inflection.
- (4) a point of local maxima.

4. A spherical iron ball of 10 cm radius is coated with a layer of ice of uniform thickness the melts at a rate of  $50 \text{ cm}^3/\text{min}$ . When the thickness of ice is 5 cm, then the rate (in cm/min.) at which of the thickness of ice decreases, is :

- (1)  $\frac{1}{36\pi}$                                   (2)  $\frac{5}{6\pi}$
- (3)  $\frac{1}{18\pi}$                                   (4)  $\frac{1}{54\pi}$

5. Let  $P(h, k)$  be a point on the curve  $y = x^2 + 7x + 2$ , nearest to the line,  $y = 3x - 3$ . Then the equation of the normal to the curve at  $P$  is :

- (1)  $x + 3y - 62 = 0$                   (2)  $x - 3y - 11 = 0$
- (3)  $x - 3y + 22 = 0$                   (4)  $x + 3y + 26 = 0$

6. If  $p(x)$  be a polynomial of degree three that has a local maximum value 8 at  $x = 1$  and a local minimum value 4 at  $x = 2$ ; then  $p(0)$  is equal to:

- (1) 12    (2) -24
- (3) 6    (4) -12

7. Suppose  $f(x)$  is a polynomial of degree four, having critical points at  $-1, 0, 1$ . If  $T = \{x \in \mathbf{R} \mid f(x) = f(0)\}$ , then the sum of squares of all the elements of  $T$  is :

- (1) 6    (2) 8
- (3) 4    (4) 2

8. If  $x = 1$  is a critical point of the function  $f(x) = (3x^2 + ax - 2 - a) e^x$ , then :

(1)  $x = 1$  is a local minima and  $x = -\frac{2}{3}$  is a local maxima of  $f$ .

(2)  $x = 1$  is a local maxima and  $x = -\frac{2}{3}$  is a local minima of  $f$ .

(3)  $x = 1$  and  $x = -\frac{2}{3}$  are local minima of  $f$ .

(4)  $x = 1$  and  $x = -\frac{2}{3}$  are local maxima of  $f$ .

9. Let  $AD$  and  $BC$  be two vertical poles at  $A$  and  $B$  respectively on a horizontal ground. If  $AD = 8 \text{ m}$ ,  $BC = 11 \text{ m}$  and  $AB = 10 \text{ m}$ ; then the distance (in meters) of a point  $M$  on  $AB$  from the point  $A$  such that  $MD^2 + MC^2$  is minimum is\_.

10. The set of all real values of  $\lambda$  for which the function  $f(x) = (1 - \cos^2 x) \cdot (\lambda + \sin x)$ ,  $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$ , has exactly one maxima and exactly one minima, is :

- (1)  $\left(-\frac{1}{2}, \frac{1}{2}\right) - \{0\}$       (2)  $\left(-\frac{1}{2}, \frac{1}{2}\right)$   
(3)  $\left(-\frac{3}{2}, \frac{3}{2}\right)$       (4)  $\left(-\frac{3}{2}, \frac{3}{2}\right) - \{0\}$

SOLUTION

1. NTA Ans. (2)


Sol.  $\lim_{x \rightarrow 0} \left( 2 + \frac{f(x)}{x^3} \right) = 4$

$\Rightarrow f(x) = 2x^3 + ax^4 + bx^5$   
 $f'(x) = 6x^2 + 4ax^3 + 5bx^4$   
 $f'(1) = 0, f'(-1) = 0$

$a = 0, b = \frac{-6}{5} \Rightarrow f(x) = 2x^3 - \frac{6}{5}x^5$

$f'(x) = 6x^2 - 6x^4$   
 $= 6x^2(1 - x)(1 + x)$

Sign scheme for  $f'(x)$



Minima at  $x = -1$

Maxima at  $x = 1$

2. NTA Ans. (3)

Sol.  $f''(x) = \lambda(x - 1)$

$f'(x) = \frac{\lambda x^2}{2} - \lambda x + C \Rightarrow f'(-1) = 0 \Rightarrow c = \frac{-3\lambda}{2}$

$f(x) = \frac{\lambda x^3}{6} - \frac{\lambda x^2}{2} - \frac{3\lambda}{2}x + d$

$f(1) = -6 \Rightarrow -11\lambda + 6d = -36 \dots(i)$

$f(-1) = 10 \Rightarrow 5\lambda + 6d = 60 \dots(ii)$

from (i) & (ii)  $\lambda = 6$  &  $d = 5$

$f(x) = x^3 - 3x^2 - 9x + 5$

Which has minima at  $x = 3$

Ans. 3.00

3. NTA Ans. (1)

Sol.  $F'(x) = x^2 g(x) = x^2 \int_1^x f(u) du \Rightarrow F'(1) = 0$

$F''(x) = x^2 f(x) - 2x \int_1^x f(u) du$

$F''(1) = 1.f(1) - 2 \times 0$

$F''(1) = 3$

$F'(1) = 0$  and  $F''(1) = 3 > 0$  So, Minima

4. NTA Ans. (3)

Sol. Let thickness of ice be 'h'.

Vol. of ice =  $v = \frac{4\pi}{3}((10+h)^3 - 10^3)$


$\frac{dv}{dt} = \frac{4\pi}{3}(3(10+h)^2) \cdot \frac{dh}{dt}$

Given  $\frac{dv}{dt} = 50 \text{ cm}^3 / \text{min}$  and  $h = 5 \text{ cm}$

$\Rightarrow 50 = \frac{4\pi}{3}(3(10+5)^2) \frac{dh}{dt}$

$\Rightarrow \frac{dh}{dt} = \frac{50}{4\pi \times 15^2} = \frac{1}{18\pi} \text{ cm / min}$

5. Official Ans. by NTA (4)



Let L be the common normal to parabola  $y = x^2 + 7x + 2$  and line  $y = 3x - 3$

$\Rightarrow$  slope of tangent of  $y = x^2 + 7x + 2$  at P = 3

$\Rightarrow \left. \frac{dy}{dx} \right|_{\text{For P}} = 3$

$\Rightarrow 2x + 7 = 3 \Rightarrow x = -2 \Rightarrow y = -8$

So P(-2, -8)


Normal at P :  $x + 3y + C = 0$

$\Rightarrow C = 26$  (P satisfies the line)

**Normal :  $x + 3y + 26 = 0$**

**6. Official Ans. by NTA (4)**

**Sol.** Since  $p(x)$  has relative extreme at



$$x = 1 \text{ \& } 2$$

so  $p'(x) = 0$  at  $x = 1$  &  $2$

$$\Rightarrow p'(x) = A(x-1)(x-2)$$

$$\Rightarrow p(x) = \int A(x^2 - 3x + 2)dx$$

$$p(x) = A\left(\frac{x^3}{3} - \frac{3x^2}{2} + 2x\right) + C \quad \dots(1)$$

$$P(1) = 8$$

From (1)

$$8 = A\left(\frac{1}{3} - \frac{3}{2} + 2\right) + C$$

$$\Rightarrow 8 = \frac{5A}{6} + C \Rightarrow \boxed{48 = 5A + 6C} \quad \dots(3)$$

$$P(2) = 4$$

$$\Rightarrow 4 = A\left(\frac{8}{3} - 6 + 4\right) + C$$

$$\Rightarrow 4 = \frac{2A}{3} + C \Rightarrow \boxed{12 = 2A + 3C} \quad \dots(4)$$

From 3 & 4,  $C = -12$

$$\text{So } P(0) = C = \boxed{-12}$$

**7. Official Ans. by NTA (3)**

**Sol.**  $f'(x) = x(x+1)(x-1) = x^3 - x$

$$\int df(x) = \int x^3 - x \, dx$$

$$f(x) = \frac{x^4}{4} - \frac{x^2}{2} + C$$

$$f(x) = f(0)$$

$$\frac{x^4}{4} - \frac{x^2}{2} = 0$$

$$x^2(x^2 - 2) = 0$$

$$x = 0, 0, \sqrt{2}, -\sqrt{2}$$

$$x_1^2 + x_2^2 + x_3^2 = 0 + 2 + 2 = 4$$

**8. Official Ans. by NTA (1)**

**Sol.**  $f(x) = (3x^2 + ax - 2 - a)e^x$

$$f'(x) = (3x^2 + ax - 2 - a)e^x + e^x(6x + a)$$

$$= e^x(3x^2 + x(6+a) - 2)$$

$$f'(x) = 0 \text{ at } x = 1$$

$$\Rightarrow 3 + (6+a) - 2 = 0$$

$$a = -7$$


$$f'(x) = e^x(3x^2 - x - 2)$$

$$= e^x(x-1)(3x+2)$$

$$\begin{array}{c} + \quad - \quad + \\ \hline -2/3 \quad 1 \end{array}$$

$x = 1$  is point of local minima

$x = \frac{-2}{3}$  is point of local maxima

**9. Official Ans. by NTA (5.00)**

**Sol.**

$$(MD)^2 + (MC)^2 = h^2 + 64 + (h-10)^2 + 121$$

$$= 2h^2 - 20h + 64 + 100 + 121$$

$$= 2(h^2 - 10h) + 285$$

$$= 2(h-5)^2 + 235$$

it is minimum if  $h = 5$

**10. Official Ans. by NTA (4)**

**Sol.**  $f(x) = (1 - \cos^2 x)(\lambda + \sin x) \quad x \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$

$$f(x) = \lambda \sin^2 x + \sin^3 x$$

$$f'(x) = 2\lambda \sin x \cos x + 3\sin^2 x \cos x$$

$$f'(x) = \sin x \cos x (2\lambda + 3\sin x)$$

$$\sin x = 0, \frac{-2\lambda}{3}, (\lambda \neq 0)$$

for exactly one maxima & minima

$$\frac{-2\lambda}{3} \in (-1, 1) \Rightarrow \lambda \in \left(\frac{-3}{2}, \frac{3}{2}\right)$$

$$\lambda \in \left(-\frac{3}{2}, \frac{3}{2}\right) - \{0\}$$