

10. Two light waves having the same wavelength λ in vacuum are in phase initially. Then the first wave travels a path L_1 through a medium of refractive index n_1 while the second wave travels a path of length L_2 through a medium of refractive index n_2 . After this the phase difference between the two waves is:

$$(1) \frac{2\pi}{\lambda}(n_1L_1 - n_2L_2) \quad (2) \frac{2\pi}{\lambda}\left(\frac{L_2}{n_1} - \frac{L_1}{n_2}\right)$$

$$(3) \frac{2\pi}{\lambda}\left(\frac{L_1}{n_1} - \frac{L_2}{n_2}\right) \quad (4) \frac{2\pi}{\lambda}(n_2L_1 - n_1L_2)$$

11. A beam of plane polarised light of large cross sectional area and uniform intensity of 3.3 Wm^{-2} falls normally on a polariser (cross sectional area $3 \times 10^{-4} \text{ m}^2$) which rotates about its axis with an angular speed of 31.4 rad/s . The energy of light passing through the polariser per revolution, is close to :

$$(1) 1.0 \times 10^{-5} \text{ J} \quad (2) 5.0 \times 10^{-4} \text{ J}$$


$$(3) 1.0 \times 10^{-4} \text{ J} \quad (4) 1.5 \times 10^{-4} \text{ J}$$

12. Orange light of wavelength $6000 \times 10^{-10} \text{ m}$ illuminates a single slit of width $0.6 \times 10^{-4} \text{ m}$. The maximum possible number of diffraction minima produced on both sides of the central maximum is _____.

13. A beam of electrons of energy E scatters from a target having atomic spacing of 1 \AA . The first maximum intensity occurs at $\theta = 60^\circ$. Then E (in eV) is _____.

(Planck constant $h = 6.64 \times 10^{-34} \text{ Js}$, $1 \text{ eV} = 1.6 \times 10^{-19} \text{ J}$, electron mass $m = 9.1 \times 10^{-31} \text{ kg}$)

14. In the figure below, P and Q are two equally intense coherent sources emitting radiation of wavelength 20 m . The separation between P and Q is 5 m and the phase of P is ahead of that of Q by 90° . A, B and C are three distinct points of observation, each equidistant from the midpoint of PQ. The intensities of radiation at A, B, C will be in the ratio:



$$(1) 0 : 1 : 2 \quad (2) 4 : 1 : 0$$

$$(3) 0 : 1 : 4 \quad (4) 2 : 1 : 0$$

15. A Young's double-slit experiment is performed using monochromatic light of wavelength λ . The intensity of light at a point on the screen, where the path difference is λ , is K units. The intensity of light at a point where the path difference is $A \frac{\lambda}{6}$ is given by $\frac{nK}{12}$, where n is an integer. The value of n is _____.

8. Official Ans. by NTA (1)

Sol. Let the length of segment is " ℓ "
Let N is the no. of fringes in " ℓ "
and w is fringe width.

→ We can write

$$N w = \ell$$

$$N \left(\frac{\lambda D}{d} \right) = \ell$$

$$\frac{N_1 \lambda_1 D}{d} = \ell$$

$$\frac{N_2 \lambda_2 D}{d} = \ell$$

$$N_1 \lambda_1 = N_2 \lambda_2$$

$$16 \times 700 = N_2 \times 400$$

$$N_2 = 28$$

9. Official Ans. by NTA (4)

Sol. $\Delta\theta_0 = \left(\frac{\lambda}{d} \times \frac{180}{\pi} \right)^0$
 $= 0.57^\circ$

10. Official Ans. by NTA (1)

Sol. $\Delta p = n_1 L_1 - n_2 L_2$

$$\Delta\phi = \frac{2\pi}{\lambda} \Delta p$$

11. Official Ans. by NTA (3)

Sol. Intensity, $I = 3.3 \text{ Wm}^{-2}$

$$\text{Area, } A = 3 \times 10^{-4} \text{ m}^2$$

Angular speed, $\omega = 31.4 \text{ rad/s}$

$$\therefore \langle \cos^2\theta \rangle = \frac{1}{2}, \text{ in one time period}$$

$$\therefore \text{Average energy} = I_0 A \times \frac{1}{2}$$

$$= \frac{(3.3)(3 \times 10^{-4})}{2}$$

$$\approx 5 \times 10^{-4} \text{ J}$$

12. Official Ans. by NTA (200)

Official Ans. by ALLEN (198)

Sol. Condition for minimum,

$$d \sin\theta = n\lambda$$

$$\therefore \sin\theta = \frac{n\lambda}{d} < 1$$

$$n < \frac{d}{\lambda} = \frac{6 \times 10^{-5}}{6 \times 10^{-7}} = 100$$


$$\therefore \text{Total number of minima on one side} \\ = 99$$

Total number of minima = 198

Correct Answer is 198

13. Official Ans. by NTA (50.00)

Sol.



$$2d \sin\theta = \lambda = \frac{h}{\sqrt{2mE}}$$

$$2 \times 10^{-10} \times \frac{\sqrt{3}}{2} = \frac{6.6 \times 10^{-34}}{\sqrt{2mE}}$$

$$E = \frac{1}{2} \times \frac{6.64^2 \times 10^{-48}}{9.1 \times 10^{-31} \times 3 \times 1.6 \times 10^{-19}} = 50.47$$

